

Prescribing Sedation for Fear of Flying

Key information




Due to a medical safety alert update from an aviation trained doctor, we have been advised that we should no longer prescribe sedating drugs called benzodiazepines (diazepam/lorazepam/temazepam or midazolam) for the use of treating fear of flying.

We have taken the decision to put our patients' safety first and to follow this safety guidance and will no longer provide prescriptions for benzodiazepines for these reasons.



The use of benzodiazepines cause longer reaction times and slowed thinking, which during a flight will put the passenger at significant risk of not being able to act in a manner which could save their life in the event of a safety critical scenario.

The use of benzodiazepines has the potential to increase the risk of DVT (deep vein thrombosis).




The sedating effects can reduce respiratory function which has the potential to be life threatening – even more so if there is a combination with alcohol.

Patients taking benzodiazepines may report that they feel increased aggression. This has the potential to put other occupants of the aircraft at risk.



Benzodiazepines are not recommended for those people with phobic states.

For some countries it is illegal to import these drugs and so the passenger will need to use a different strategy for the homeward bound journey and/or any subsequent legs of the journey.



We recommend patients who have fear of flying to research going on fear of flying courses. These are run by several major airlines and sometimes by local airports. We are unfortunately unable to recommend any specific courses

If you would like to discuss any of these matters further, please call the surgery for an appointment on 01422 415777.

