Plantar Fasciitis Exercises

London North West Healthcare NHS Trust

http://www.lnwh.nhs.uk/services/msk-exercises

GP Generic Ú|æ) æk Á Øæe & ãitã Exercises Tæî Á 2011

Stretch gently. Stop any exercise causing tingling or severe pain.

- Sit on the floor with your legs stretched out in front
- Loop a towel around the ball of your foot and pull the towel stretching the back of your calf
- Hold for 30 seconds. Repeat 3 times daily

© PhysioTools Ltd



- Stand holding onto a stable support such as a chair or the wall with your hands on the support
- Keep the sore foot back and your good foot forward, and the heel of your injured foot firmly on the floor

- Turn your injured (back) foot slightly inward and slowly lean into the wall until you feel a stretch in the back of your calf

- Hold for 30 seconds. Repeat 3 times daily

© PhysioTools Ltd



- Stand holding onto a stable support such as a chair or the wall with your hands on the support- Keep your knees a little bent and the sore foot back
- Gently lean your knees in towards the chair to stretch the back of the calf
- Keep your heel on the ground
- Hold for 30 seconds and repeat 3 times daily

© PhysioTools Ltd



- Place the affected foot over the opposite knee
- Place your fingers around the base of the toes and pull back towards the shin until you can feel a stretch in the arch (plantar fascia)
- Hold this position for 30 seconds, ropest 3 times daily
- Hold this position for 30 seconds, repeat 3 times daily