Special Considerations for premature babies being discharged from Neonatal Units

Your baby may well behave differently once you are home and cry more from 2 weeks after the date they were due to be born.

- Premature babies are prone to crying more often and can prove difficult to soothe. This phase will improve over time.
- Following discharge some babies find it difficult to adjust to the quiet of home but this will get better.
- You will soon learn how much crying is normal for your baby but if you think there is something wrong with your baby then you must seek medical help.

Share the ICON message!

It isn't just parents who get frustrated at a baby's cry. Think very carefully about who you ask to look after your baby.

Share the ICON message with anyone who may look after your baby.

Check that caregivers understand about how to cope with crying before you decide to leave your baby with them and share this ICON leaflet with them.

Reminder about Safe Sleeping:

You can talk to your Midwife or Health Visitor about all aspects of crying and safe sleeping.

Further information and support

Health Visiting Services

ChatHealth is a text messaging service which supports parents, carers and families of children under 5 in Somerset. There is also an extended service for older children. How to use ChaHealth: Text the dedicated phone number: **07480635514**

Midwifery Services

Musgrove Park Hospital Bracken Birth Centre 01823 342058
Veovil District Hospital Freya Ward

Yeovil District Hospital Freya Ward 01935 384303

Somerset Childrens Social Care.
If you are worried about the safety of a child, call 0300 123 2224

NHS Choices

The NHS Webpage has information on how to soothe a crying baby which can be accessed via www.nhs.uk Wessex Healthier Together www.what0-18.nhs.uk

National Society for the Prevention of Cruelty to Children (NSPCC) Tel: 0808 800 5000 www.nspcc.org.uk

For more information on ICON and aspects of maternity and parenting, visit: www.cypsomersethealth.org

CRY-SIS National Help Line: 08451 228669
Lines open 7 days a week, 9am-10pm
Remember – if you are concerned that your baby may
be unwell, contact your GP or NHS 111
(go to 111.nhs.uk or call 111- the service
is available 24 hours a day, 7 days a week).
In an emergency, ring 999.



Twitter: ICON_COPE

For a translation of this document, an interpreter or a version in

rge or Braille









Infant crying and how to cope

Babies born prematurely



Information for parents and carers taking premature babies home





Babies Cry!

Infant crying is normal and it will stop.


A baby's cry can be upsetting and frustrating. It is designed to get your attention and you may be worried that something is wrong with your baby.

Your baby may start to cry more frequently at about 2 weeks of age. The crying may get more frequent and last longer during the next few weeks, hitting a peak at about 6 to 8 weeks.

Every baby is different, but after about 8 weeks, babies start to cry less and less each week.

Normal Crying Curve Curves of Early Infant Crying

2 weeks to 4-5 months



Barr, R. (1990) The normal crying curve: What do we really know?

Developmental Medicine and Child Neurology.

What can I do to help my baby?

Comfort methods can sometimes soothe the baby and the crying will stop.

Babies can cry for reasons such as if they are hungry, tired, wet/dirty or if they are unwell.

Check these basic needs and try some simple calming techniques:

- Talk calmly, hum or sing to your baby
- Let them hear a repeating or soothing sound
- Hold them close skin to skin
- Go for a walk outside with your baby
- Give them a warm bath

These techniques may not always work. It may take a combination or more than one attempt to soothe your baby.

If you think there is something wrong with your baby or the crying won't stop speak to your GP, Midwife or Health Visitor. If you are worried that your baby is unwell call NHS 111.

The crying won't stop, what can I do now?

Not every baby is easy to calm but that doesn't mean you are doing anything wrong.

Don't get angry with your baby or yourself. Instead, put your baby in a safe place and walk away so that you can calm yourself down by doing something that takes your mind off the crying. Try:

- Listening to music, doing some exercises or doing something that calms you.
- Call a relative or friend they may be able to help you calm or may be able to watch your baby.

After a few minutes when you are calm, go back and check on the baby.

It's normal for parents to get stressed, especially by crying. Put some time aside for yourself and take care of your needs as well as your baby's to help you cope.

What not to do...

Handling a baby roughly will make them more upset. Shouting or getting angry with your baby will make things worse.

Sometimes parents and people looking after babies get so angry and frustrated with a baby's cry they lose control.

They act on impulse and shake their baby.

Shaking or losing your temper with a baby is very dangerous and can cause:

- Blindness
- Learning disabilities
- Seizures
- Physical disabilities
- Death

Remember: Never ever shake or hurt a baby



Remember – This phase will stop! Be an ICON for your baby and cope with their crying.

- Infant crying is normal and it will stop
- Comfort methods can sometimes soothe the baby and the crying will stop
- O It's OK to walk away if you have checked the baby is safe and the crying is getting to you
- Never ever shake or hurt a baby