

## PRECRIBING MOUNJARO (TIRZEPATIDE) IN GENERAL PRACTICE FOR WEIGHT LOSS

## AT PRESENT, AND UNTIL AT LEAST **OCTOBER 2025**, GENERAL PRACTITIONERS **CANNOT** PRESCRIBE MOUNJARO (TIRZEPATIDE) FOR WEIGHT LOSS

Although NICE issued guidance in December 2024 that the NHS should provide Mounjaro (tirzepatide) to patients for weight loss from June 2025, presently it is unknown how NHS Leicester, Leicestershire and Rutland (NHS LLR) will be arranging for it to be given. The service will not start until at least October 2025 (and possibly later).

NHS England has provided funding for a very limited number of patients to be prescribed tirzepatide (425 in total for this year for the whole of Leicester, Leicestershire and Rutland).

NHS LLR has also decided that patients cannot be referred for NHS care to Right to Choose providers like 'Oviva.'

The criteria are very strict (see below - for this year a BMI greater than 40, and at least four qualifying medical problems), but even so the number of eligible patients in LLR will be far in excess of the number being funded.

Patients with any further questions or concerns can contact NHS LLR:

Room G30, Pen Lloyd Building, County Hall, Glenfield, Leicester, LE3 8TB Email: llricb-llr.enguiries@nhs.net Telephone: 0116 295 3405

Year	<b>Estimated Cohort Duration</b>	Cohort	Cohort Access Groups	
			Comorbidities	BMI*
Year 1	12 months	1	≥4 'qualifying' comorbidities:	≥ 40
2025/6			<ul> <li>Hypertension</li> </ul>	
			<ul> <li>Dyslipidaemia</li> </ul>	
			Obstructive Sleep Apnoea	
			Cardiovascular disease	
			Type 2 diabetes mellitus	
Year 2 2026/7	9 months	П	≥4 'qualifying' comorbidities:	35 – 39.9
			<ul> <li>Hypertension</li> </ul>	
			<ul> <li>Dyslipidaemia</li> </ul>	
			Obstructive Sleep Apnoea	
			Cardiovascular disease	
			Type 2 diabetes mellitus	
Year 2/3	15 months	Ш	3 'qualifying' comorbidities:	≥ 40
2026 and			<ul> <li>Hypertension</li> </ul>	
2027/8			<ul> <li>Dyslipidaemia</li> </ul>	
			Obstructive Sleep Apnoea	
			Cardiovascular disease	
			Type 2 diabetes mellitus	

<sup>\*</sup>Use a lower BMI threshold (usually reduced by 2.5 kg/m2) for people from South Asian, Chinese, other Asian, Middle Eastern, Black African or African-Caribbean ethnic backgrounds.

**Version 3. Information Accurate as of 27 August 2025** 

## ADVISE SUPPORT REPRESENT

Leicester, Leicestershire and Rutland Local Medical Committee 109 Station Road, Glenfield, Leicester, LE3 8GS enquiries@llrlmc.co.uk www.llrlmc.co.uk 0116 296 2950