

Eyes Right

Helping non-ophthalmic
health care professionals
deliver quality care safely



Who is this guide for?

All non-ophthalmic nursing staff, including health care assistants, in adult care settings ranging from non-ophthalmic hospital settings to residential and nursing homes. This guide has been developed following a review of the research literature and in conjunction with the expert advice of the RCN Ophthalmic Forum members and advisers.

This document has been designed in collaboration with our members to ensure it meets most accessibility standards.

However, if this does not fit your requirements, please contact **corporate.communications@rcn.org.uk**

RCN Legal Disclaimer

This publication contains information, advice and guidance to help members of the RCN. It is intended for use within the UK but readers are advised that practices may vary in each country and outside the UK. The information in this booklet has been compiled from professional sources, but its accuracy is not guaranteed. Whilst every effort has been made to ensure the RCN provides accurate and expert information and guidance, it is impossible to predict all the circumstances in which it may be used. Accordingly, the RCN shall not be liable to any person or entity with respect to any loss or damage

caused or alleged to be caused directly or indirectly by what is contained in or left out of this website information and guidance.

Published by the Royal College of Nursing, 20 Cavendish Square, London W1G 0RN

© 2022 Royal College of Nursing. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means electronic, mechanical, photocopying, recording or otherwise, without prior permission of the Publishers. This publication may not be lent, resold, hired out or otherwise disposed of by ways of trade in any form of binding or cover other than that in which it is published, without the prior consent of the Publishers.



Contents

- 03** How will this guide help me
- 03** Supporting the visually impaired person
- 04** Advice about regular eye tests
- 05** Eye drop and eye ointment instillation
- 07** Cleaning the eye
- 09** Daily living skills
- 10** Mobility
- 11** Ophthalmic emergencies
- 13** Non-emergency conditions requiring attention
- 14** Accessing services





How will this guide help me?

This guide will highlight the importance of eye care, the role and responsibility of the carer in delivering care and accessing the right care at the right time. Please also refer to local policies and procedures, where applicable.

The guide has focused on the following topics:

- supporting the visually impaired person
- eye drop and eye ointment instillation
- cleaning the eye.

Ophthalmic emergencies:

- red eye
- flashing lights and floaters
- ocular pain
- copious discharge
- foreign body sensation
- chemical injury.

Non-emergency conditions requiring attention:

- entropion
- ectropion
- blepharitis.



Supporting the visually impaired person

Prevalence of visual impairment

- There are over two million people in the UK living with sight loss.
- One in five people aged 75 and over are living with sight loss.
- One in two people aged 90 and over are living with sight loss.

Adults with learning disabilities are 10 times more likely to have serious sight problems than other adults and children (Public Health England, 2020).

Advice about regular eye tests

Is the person wearing the correct spectacles?

An older person may get confused with their spectacles if they have two pairs and may attempt to watch television with reading spectacles on.

Has the person had a recent eye test?

Sight tests should be undertaken on a yearly basis unless there is a change in vision, in which case an earlier appointment should be made.

If the service user has received notification from either an ophthalmic clinic or optician, that their vision loss falls into one of these three categories, they may be suitable for registration as visually impaired:

- visual acuity of 3/60 to 6/60 with full field of vision
- visual acuity of up to 6/24 with moderate reduction of visual field plus central vision is cloudy or blurred
- visual acuity of 6/18 or even better if half or more of the visual field is missing.

Common eye conditions may affect vision, for example:

Cataracts
Causes blurring of vision and can affect contrast sensitivity.

Age-related macular degeneration
Can cause visual distortion and affect the central vision.

Glaucoma
Can affect both central and peripheral field of vision.

Certification of visual impairment (CVI) may be applied for in clinic by a consultant ophthalmologist, in order for people to obtain support with daily living and maintain independence.



Eye drop and eye ointment instillation

- Gather your equipment, including the drops, prescription, tissues or cotton wool balls.
- Store the eye drops and ointment as per manufacturer's instructions. Wear gloves, as required, for example if likely to be in direct contact with the conjunctiva or if there is a risk of contamination from eye lids or orbital wounds.

Aprons are not required routinely unless there is a risk of gross contamination of uniform.
- Wash or gel your hands and allow to dry thoroughly before and after instilling the drops or if the hands become contaminated during the procedure. Gloves are required if hands are in contact with the conjunctiva, or if there is a risk of contamination from eye lids or orbital wounds.
- If the patient has had eye surgery in the last 24-48 hours, the procedure should be aseptic.
- Prepare your patient (sitting or lying with the head supported) and obtain consent.
- Establish you have identified the correct eye.
- If contact lenses are worn, these should be removed unless instructed otherwise, and re-inserted after 30 mins, if trained and assessed to do so, as per local policy.
- Examine the eye(s) before instilling the drops/ointment to look for signs of improvement or deterioration or allergy.
- You may need to clean the eye lids if required, for example if there is crusting on the eye lids.
- Check that the drops/ointment have not expired and are due to be instilled. Gently agitate the bottle as settling may have occurred, if appropriate.
- Remove the cap and place it on a clean surface.
- Warn the patient that the eye drops may sting or ointment will cause blurring of vision – this is temporary.

- Ask the patient to look up. Using the tissue, gently pull down the lower eye lid to form a pocket (fornix) in which to instil the eye drop/apply the ointment. Instil the drop into the fornix or apply a ribbon of ointment.
- Release the eye lid and ask the patient to gently close their eye for a slow count of 60. This will help the drug to absorb.
- Allow the patient to mop their cheek to clear excess drop and for comfort, or wipe excess ointment from the lids.
- If another eye drop is required, wait five minutes between eye drops to the same eye.
- If ointment is required, always instil the eye drop first.
- Make the patient comfortable.
- Clean and replace the patient's spectacles, if worn.
- Complete your documentation.
- Take every opportunity to teach the patient or carer to instil their own eye drops or apply their ointment.



Right...

...patient

...medication

...dose

...route

...time

...to refuse

...assessment

...evaluation


...documentation

...patient education



Cleaning the eye

- Identify the patient and explain the procedure to obtain consent.
- Make them comfortable, sitting or lying, with the head supported.
- Wash or gel the hands before and after the procedure or if the hands become contaminated during the procedure.
- If the eye lids have been operated on and sutures are present, use non-linting and sterile swabs to clean the eyelids, otherwise use sterile cotton wool balls. Follow post-operative instructions as advised.
- In the home, use cooled boiled water, contained in a clean receptacle. In hospital or residential nursing or care settings, use sachets of sterile normal saline or sterile water.
- A specialist eye dressing pack may be used. If not a sterile dressing pack may be used, but do not use any forceps as they may damage the eye or eye lids. The pack should be placed on a clean surface..
- Protect the patient's clothing with a drape, such as a clean tea towel.
- Examine the eye lids and eye with a clean pen-torch and note and report any abnormalities such as redness, increased temperature, unusual swelling or presence of discharge.
- Ask the patient to close their eye lids.
- Moisten the swab using one hand. Transfer the swab to the other hand. Gently clean the eyelids from the nasal side to the outer corner.
- Discard the swab into a bag/ container away from the sterile or clean materials.

- 
- Take another swab as before. Ask the patient to look up so as to avoid damaging the cornea. Swab the lower lid, taking care to clear debris from the lashes.
 - Repeat as required until the lids are clean. Be careful not to make the lids sore.
 - Now take a dry swab, ask the patient to close their lids and swab from the inner to outer edge of the lids to dry them.
 - Make the patient comfortable.
 - Clean and replace the patient's spectacles, if worn.
 - Clear away the equipment.
 - Document your findings and inform your patient.





Daily living skills

Assess the environment for:

- lighting – it is important to check the correct lighting is in use. For example, energy saving light bulbs have a delay before illumination, so fluorescent lights are better to prevent slips trips and falls.
- safety – hazards around the home should be identified and risks reduced, eg. hand rails and support aids provided.

What can help?

Low vision aids – can incorporate new technology, eg. mobile phone or tablets can be used to increase font size or adjust contrast.

Magnifiers – are also used to enlarge print for reading. The principles are to make things

bigger, brighter, bolder!

Liquid level indicators can be used to make a hot drink safely and sight awareness guide training by local authorities can be arranged.



- Increased awareness aims to reduce the risk of falls.
- Cane training arranged by local authorities or sight loss organisations will help enable independence.
- Guide dog application can be considered.
- Learned routes so people gain knowledge of common routes they need to use as part of everyday living.
- In-home mobility training may be required for sudden visual loss, to familiarise people with potential risks to safety in the home.
- If help is needed with technology the following link may be useful - pocklington-trust.org.uk/supporting-you/useful-guides/technology-guides-for-you



Ophthalmic emergencies

In the event that you encounter any of the following circumstances:

- call NHS 111
- attend the nearest A&E Department.

Signs and symptoms

- Sudden increase in ocular pain.
- Sudden loss of vision rather than gradual change in vision.
- Bleeding in or around the eye.
- Photophobia/sensitivity to light.
- Redness and abrasion.
- Ocular pain with headache and abdominal pain.
- Sudden painful swelling of the upper lid with an accompanying fever and feeling unwell.

Common emergency conditions

Red eye

- The patient has a painful red eye and has other symptoms, such as any change of vision, sensitivity to light, a severe headache and/or flashing lights.
- The patient has recently injured their eye.
- New or excessive amounts of mucus, especially green, present with changes in vision or eye pain requires medical evaluation to determine the underlying cause.




Fig. 1

Flashing lights and floaters

- A new onset of floaters and flashing light in the eye.
- Gradual shading of vision from one side, like a curtain being drawn.
- Rapid decline in sharp central vision.

Foreign body sensation

- A foreign body is an object in the eye that shouldn't be there, such as a speck of dust, a wood chip, a metal splinter, an insect or a piece of glass.
- Don't remove a foreign body or attempt to remove it.




Fig. 2

Chemical injury

- Chemical splashes, such as from corrosive substances, into the eye is an emergency as damage occurs quickly.
- The patient may experience and report a burning sensation, excessive tearing, pain, redness in the eyelid surface or blurred vision.
- Irrigate the eye(s) immediately with copious amounts of cold water.
- The nurse is reminded to always use their professional judgment, when assessing urgent care needs, and if concerned or in the event of an emergency, consider dialing 999 for emergency care.




Fig. 3

Figs. 1, 2, 5, 6, 7 = Figs. 4.9, 5.3, 1.11, 4.5, 4.6 by permission from Jackson TL; Moorfields Manual of Ophthalmology 2nd Edition; JP Medical Ltd 2014.

Fig. 3 = Fig. 4.22 Courtesy of Mr. DH Verity.

Fig. 4 = Fig. 1.10 Courtesy of Mrs. N Dunlop.

Non-emergency conditions requiring attention

In the event that you encounter any of the following circumstances, please contact the patient's Optometrist or GP.

Entropion

- Either the upper and/or lower eyelid may turn inwards onto the eye, causing irritation, pain and watering of the eye.
- Untreated, the lashes rub on the cornea, damaging it and therefore affecting vision.



Fig. 4

Ectropion

- The lower eye lid droops away from the eye to the point where you may see the pink inner surface of the eyelid.
- Untreated, several structures may be affected:
 - cornea – may become dry and sore, which may lead to a corneal ulcer
 - tear duct – may be displaced so tears drain down the face instead of into the tear duct
 - dry eye – as tears are not staying long enough on the surface of the eye to lubricate the eye, more tears are produced to remedy this. This leads to a constantly watery eye.



Fig. 5

Blepharitis

- Chronic inflammation of the eyelids can leave the lids sore and often crusted.
- Treatment is normally eyelid hygiene as discussed in the section on Cleaning the eye.



Fig. 6




Fig. 7





Accessing services

Accessing services

Help can be sought in the following ways:

- medical referral from the GP
- for visually impaired individuals via CVI England, Wales and Scotland and A655 Northern Ireland registration: it is standard practice that a copy of the registration form, completed by the Ophthalmologist, will be sent to the person's local social services department
- self-referral can also be made to the local authority services.

Useful contact numbers

RNIB helpline 0303 1239999 or rnib.org.uk

A 'buddy system' can be arranged, whereby people are put in contact with a visually impaired person who is coping positively with visual loss.

Patient support groups include

Macular Society: 0300 3030111 or help@macularsociety.org

Glaucoma UK: 01233 648170 or helpline@glaucoma.uk

For any of the listed common ophthalmic emergencies, please seek medical assistance. If you are unsure, please seek medical assistance.



Royal College
of Nursing

Publication

This is an RCN practice guidance. Practice guidance are evidence-based consensus documents, used to guide decisions about appropriate care of an individual, family or population in a specific context.

Publication description

This updated guide highlights the importance of eye care, the role and responsibilities of the carer in delivering care and accessing the right care at the right time. It has been developed for all non-ophthalmic nursing staff in adult care settings, ranging from non-ophthalmic hospital settings to residential and nursing homes.

Publication date: July 2022

Review date: November 2024

The Nine Quality Standards

These guidelines do not replace clinical assessment and any personal health queries or concerns should always be directed to an appropriate health care professional.

This publication has met the nine quality standards of the quality framework for RCN professional publications. For more information, or to request further details on how the nine quality standards have been met in relation to this particular professional publication, please contact publications.feedback@rcn.org.uk

Evaluation

The authors would value any feedback you have about this publication. Please contact publications.feedback@rcn.org.uk clearly stating which publication you are commenting on.