



BRETTON PARK HEALTHCARE

the service we provide is the service we are happy to receive

Address: Bretton Park Healthcare, Rightwell East, Bretton, Peterborough. PE3 8DT

Telephone: 01733264506

Website: www.brettonparkhealthcare.co.uk

Information for patients with type 2 diabetes on SGLT2 inhibitors.

This leaflet provides information regarding benefits, possible side effects including DKA and circumstances in which patients need to seek medical advice.

What are SGLT2 inhibitors?

- SGLT2 inhibitors are a class of drugs used in the treatment of diabetes.
- They are available in tablet form and usually taken once daily.
- They may be prescribed on their own or in combination with other diabetes medications including insulin.
- They help lower blood glucose level by re-ducing the reabsorption of glucose in the kidneys and allowing glucose to be passed out in the urine.

Who are SGLT2 inhibitors suitable for? This medication is used for the treatment of type 2 diabetes. Before starting this medication tell your doctor/diabetes specialist nurse if you have the following conditions:

- Kidney disease.
- Previous history of diabetic ketoacidosis (DKA).
- Problems with recurrent urinary tract infections.
- Problems with recurrent genital infections. ('thrush').
- History of peripheral vascular disease.
- Alcohol dependency.
- Planning or are pregnant or breastfeeding.

If you have any of the above conditions, SGLT2 inhibitor may not be suitable for you.

What are the benefits of SGLT2 inhibitors?

- Improvement in blood glucose.
- Weight loss.
- Lowering of blood pressure.
- Lowering risks of death in people with heart disease.

What are the possible side effects of SGLT2 inhibitors?

Very common ($\geq 1/10$)

An increase in the risk of hypoglycaemia when used in combination with other diabetes medication (insulin/sulfonylurea).

Common ($\geq 1/100$ to $< 1/10$)

- Developing genital or urinary tract infections.
- Passing more urine more often.
- Increased thirst.
- Itching/rash.

Uncommon ($\geq 1/1,000$ to $< 1/100$)

- Getting low blood pressure and dehydration.
- Difficulty passing urine (dysuria).

Rare ($\geq 1/10,000$ to $< 1/1,000$)

- Diabetic ketoacidosis (*rare but serious side effect*).
- Necrotising fasciitis of the perineum (Fournier's gangrene) [pain, tenderness, erythema, or swelling in the genital or perineal area, with fever or malaise].

What is diabetic ketoacidosis (DKA)?

- DKA is rare but serious condition that can develop in people taking SGLT 2 inhibitor.

It is a serious condition which requires urgent medical treatment.

- DKA usually occurs when the body does not have enough insulin, this result in the formation of ketone bodies leading to increasing levels of acid in the blood.

What are the symptoms of DKA?

- Nausea and vomiting.
- Abdominal/stomach pain.
- Rapid breathing.
- Dehydration e.g. dizziness and excessive thirst, a sweet or metallic taste in the mouth or a different odour to urine or sweat.
- Drowsiness or tiredness.
- Confusion.
- In some cases of DKA blood glucose levels maybe in the normal range in people using SGLT2 inhibitors.

If you experience any of these symptoms even if your blood glucose levels are not high, you should contact your doctor or diabetes nurse or go to the nearest hospital and tell the emergency doctor that you are taking a SGLT2 inhibitor.

See your doctor urgently:

- If you have a severe illness.
- If you are very dehydrated.
- If you have had significant weight loss and have poor appetite.
- If there is excess alcohol use.
- If you are on insulin treatment and there is a sudden reduction in insulin doses.

Stop taking your SGLT2 inhibitor in these circumstances:

- At least three days before fasting for surgery or a procedure.
- If you are unable to eat or drink e.g. nausea, vomiting.
- If you are unwell with an infection or illness.
- If you are on a very low carbohydrate diet.

SGLT2 inhibitors available in Europe:

Canagliflozin

- Invokana (Canagliflozin)
- Vokanamet (Canagliflozin / Metformin)

Dapagliflozin

- Ebymect (Dapagliflozin/metformin)
- Edistride (Dapagliflozin)
- Forxiga (Dapagliflozin)
- Qtern (Dapagliflozin/Saxagliptin)
- Xigduo (Dapagliflozin / Metformin)

Empagliflozin

- Jardiance (Empagliflozin)
- Synjardy (Empagliflozin / Metformin)

Ertugliflozin

- Steglatro (Ertugliflozin)